## AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

- (Canceled) l. 2. (Canceled) (Canceled) 3. (Canceled) 4. 5. (Canceled) 6. (Canceled) (Canceled) 7. 8. (Canceled) (Currently Amended) A composition according to claim [[1]] 19, wherein said 9. composition has a pH from about 8 to about 12. (Currently Amended) A composition according to claim [[1]] 19, wherein said 10. composition is in the form of an oil-in-water emulsion. (Currently Amended) A composition according to claim [[1]] 19, wherein said 11. composition is in the form of a thickened aqueous solution.
  - Page 3 of 8

(Canceled)

12.

- 13. (Canceled)
- 14. (Canceled)
- 15. (Canceled)
- 16. (Canceled)
- 17. (Canceled)
- 18. (Canceled)
- 19. (Currently Amended) A composition suitable for treating hair comprising:
  - a) an oxidizing agent; and
  - b) a chelant (L) having a  $\frac{\log K_{\text{CuL}}}{\log K_{\text{Cal}}}$  ratio calculated at pH 10 of at least about 3.20, wherein  $\log K_{\text{CuL}}$  is the common logarithm of the Conditional Stability Constant of said chelant with  $\text{Cu}^{2+}$  and  $\log K_{\text{CaL}}$  is the common logarithm of the Conditional Stability Constant of said chelant with  $\text{Ca}^{2+}$ :

wherein said chelant is in an amount sufficient to provide a Normalized Shinc Ratio of at least about 0.95 as measured by the Goniophotometer Damage Assessing Protocol after a 5-Cycle Hair Oxidative Treatment Protocol With 10 Intermediate Washes.

- 20. (Canceled)
- 21. (Currently Amended) A composition according to claim [[20]] 19, wherein said chelant (L) has a Hydrogen Peroxide Decomposition Ratio (% Loss) of less than about 10% as measured by the Hydrogen Peroxide Decomposition Ratio Measurement Protocol.

- (Currently Amended) A composition according to claim [[20]] 19, wherein said chelant (L) is capable of forming a hexadendate complex with Cu<sup>2+</sup>.
- 23. (Currently Amended) A composition according to claim [[20]] 19, wherein said chelant (L) is an aminocarboxylic acid chelant selected from the group consisting of diamine-N,N'-dipolyacids, monoamine monoamide-N,N'-dipolyacids and N,N'-bis(2-hydroxybenzyl)cthylenediamine-N,N'-diacetic acid (HBED), salts thereof, derivatives thereof and mixtures thereof.
- 24. (Original) A composition according to claim 23, wherein said aminocarboxylic acid chelant comprises at least two acid groups independently selected from a carboxylic acid group (-COOH), a sulphonic group (-SO<sub>3</sub>H), an o-hydroxyphenyl group, the m-hydroxyphenyl group, and a p-hydroxyphenyl group.
- 25. (Original) A composition according to claim 24, wherein said aminocarboxylic acid chelant is selected from the group consisting of ethylenediamine-N,N'-disuccinic acid (EDDS), ethylenediamine-N,N'-diglutaric acid (EDDG), 2-hydroxypropylenediamine-N,N'-disuccinic acid (HPDDS), glycinamide-N,N'-disuccinic acid (GADS), ethylenediamine-N-N'-bis(ortho-hydroxyphenyl acetic acid) (EDDHA), and salts thereof, derivatives thereof and mixtures thereof.
- 26. (Original) A composition according to claim 19, wherein said oxidizing agent is present at a level of from about 0.1% to about 40% by weight of the composition and is selected from water-soluble oxidizing agents and mixtures thereof.
- 27. (Original) A composition according to claim 19, further comprising at least one oxidative hair dye precursor.
- 28. (Currently Amended) A method of treating hair, said method comprising the steps of:
  - i) applying a first composition comprising an oxidizing agent;

JAN-02-2007

applying a second composition comprising a chelant (L) having a ii) log K<sub>CuL</sub> ratio calculated at pH 10 of at least about 3.20, wherein log K<sub>CuL</sub> is the log K Cat. common logarithm of the Conditional Stability Constant of said chelant with Cu2+ and log K<sub>CaL</sub> is the common logarithm of the Conditional Stability Constant of said chelant with Ca2+;

wherein said chelant is in an amount sufficient to provide a Normalized Shine Ratio of at least about 0.95 as measured by the Goniophotometer Damage Assessing Protocol after a 5-Cycle Hair Oxidative Treatment Protocol With 10 Intermediate Washes; and

- applying a third composition comprising a second oxidizing agent; and iii) wherein steps i) and iii) are separated by at least 1 day and step ii) does not take place immediately before step iii).
- (Original) A kit for dyeing hair comprising a first and a second compositions 29. packaged in different containers, wherein said first composition comprises an oxidizing agent and said second composition comprises an oxidative dye precursor, wherein the resulting mixture of said first and second compositions is a composition according to claim 27.
- (Original) A method of dycing human hair, said method comprising the steps of: 30.
  - i) mixing the first and second composition of a kit according to claim 29;
  - ii) applying the mixture obtained after step i) to hair;
  - iii) massaging said mixture into hair;
  - iv) retaining said mixture on the hair for an amount of time sufficient for mixture to dye the hair;
  - iv) rinsing off said composition with water.
- (New) A method of treating hair, said method comprising the steps of: 31.
  - applying to the hair a first composition comprising a chelant (L) having a i)  $\frac{\log K_{Cul}}{2}$  ratio calculated at pH 10 of at least about 3.20, wherein log

P.08/09

Atty. Dacket No. CM2517M2CC Customer No. 27752

 $K_{CuL}$  is the common logarithm of the Conditional Stability Constant of said chelant with  $Cu^{2+}$  and log  $K_{CaL}$  is the common logarithm of the Conditional Stability Constant of said chelant with  $Ca^{2+}$ ;

wherein said chelant is in an amount sufficient to provide a Normalized Shine Ratio of at least about 0.95 as measured by the Goniophotometer Damage Assessing Protocol after a 5-Cycle Hair Oxidative Treatment Protocol With 10 Intermediate Washes; and

- ii) applying to the hair a second composition comprising an oxidizing agent; wherein step i) takes place before step ii).
- (New) A method of treating hair according to claim 31, wherein step i) takes place immediately before step ii).
- 33. (New) A method of treating hair according to claim 31, wherein said first composition is not rinsed off the hair before said second composition is applied to the hair.
- 34. (New) A kit for treating hair comprising:
  - i) a first separately packaged composition comprising a chelant (L) having a  $\frac{\log K_{\text{cut}}}{\log K_{\text{cat}}}$  ratio calculated at pH 10 of at least about 3.20, wherein  $\log K_{\text{CuL}}$  is the common logarithm of the Conditional Stability Constant of said chelant with  $\text{Cu}^{2+}$  and  $\log K_{\text{CaL}}$  is the common logarithm of the Conditional Stability Constant of said chelant with  $\text{Ca}^{2+}$ ;

wherein said chelant is in an amount sufficient to provide a Normalized Shine Ratio of at least about 0.95 as measured by the Goniophotometer Damage Assessing Protocol after a 5-Cycle Hair Oxidative Treatment Protocol With 10 Intermediate Washes; and

ii) a second separately packaged composition comprising an oxidizing agent.

Please direct further correspondence to: Customer No. 27752

Respectfully submitted,

The Procter & Gamble Company

Marianne Dressman Registration No. 42,498

(513) 626-0673

Date: January 2, 2007

Customer No. 27752